



# Screening, Referral and Treatment for Attention Deficit and Hyperactivity Disorder (ADHD) – Adult – Ambulatory Clinical Practice Guideline

## Table of Contents

<b>EXECUTIVE SUMMARY</b> .....	<b>3</b>
<b>SCOPE</b> .....	<b>4</b>
<b>METHODOLOGY</b> .....	<b>5</b>
<b>INTRODUCTION</b> .....	<b>5</b>
<b>RECOMMENDATIONS</b> .....	<b>6</b>
1. PRESENTATION AND SCREENING .....	7
2. CLINICAL ASSESSMENT .....	7
3. ESTABLISH DIAGNOSIS.....	9
4. PROVIDE TREATMENT .....	11
5. COMPLETE FOLLOW-UP CARE .....	12
<b>UW HEALTH IMPLEMENTATION</b> .....	<b>13</b>
<b>REFERENCES</b> .....	<b>14</b>
<b>APPENDIX A</b> .....	<b>15</b>
<b>APPENDIX B</b> .....	<b>16</b>
<b>APPENDIX C</b> .....	<b>17</b>

**Note: Active Table of Contents -- Click to follow link**

**CPG Contact for Content:**  
Name: Peggy Scallon, MD - Psychiatry  
Phone Number: (608) 263-6100  
Email Address: [pscallon@wisc.edu](mailto:pscallon@wisc.edu)

**CPG Contact for Changes:**

Name: Lindsey Spencer, MS – Center for Clinical Knowledge Management (CCKM)

Phone Number: (608) 890-6403

Email Address: [lspencer2@uwhealth.org](mailto:lspencer2@uwhealth.org)

**Coordinating Team Members:**

Greg Rogers, PhD – Psychology

Julie Gocey, MD - Pediatrics

Nicole Weathers, MD – Family Medicine

Lisa Simpson, PA – Family Medicine

Maureen Van Dinter, NP – Family Medicine

Henny Regnier, NP – Family Medicine

Cindy Gaston, PharmD – Drug Policy Program

MaryAnn Steiner, PharmD – Unity Health Insurance

Kim Hein-Beardsley – Unity Health Insurance

Lori Samolyk – Physicians Plus Insurance Corporation

Katherine Schmitt, MD – Meriter

**Committee Approvals/Dates:**

Clinical Knowledge Management (CKM) Council (10/23/2014)

**Release Date:** October 2014

**Next Review Date:** October 2016

# Executive Summary

## Guideline Overview

This document has been developed to assist in identifying, treating, and monitoring adult patients with potential or diagnosed ADHD.

## Key Practice Recommendations

1. Assess symptoms and functional impairment
2. Complete physical exam and consider comorbid or alternative diagnoses
3. Establish ADHD diagnosis using DSM-5 diagnostic criteria
4. Provide behavioral and/or pharmacotherapy
5. Perform periodic follow-up to confirm treatment efficacy and absence of side effects

## Companion Documents

1. [Adult ADHD Algorithm](#)
2. [Adult ADHD Medication Algorithm](#)
3. [Adult Medication Charts](#)
4. [Adult ADHD Self-Report Scale \(ASRS-v1.1\) Symptom Checklist](#)

### Related Guidelines:

1. [UW Health Alcohol – Pediatric/Adult – Ambulatory Guideline](#)
2. [UW Health Tobacco – Pediatric/Adult – Inpatient/Ambulatory Guideline](#)
3. [UW Health Depression – Pediatric/Adult – Ambulatory Guideline](#)

### External Resources

1. [Wisconsin Prescription Drug Monitoring Program \(PDMP\)](#)
2. [Wisconsin Uniform Controlled Substances Act](#)

## Patient Resources

1. Healthwise: ADHD (Attention Deficit Hyperactivity Disorder): Adult
2. Healthwise: ADHD: Adults: General Info
3. [Health Information: ADHD \(Attention Deficit/Hyperactivity Disorder\)](#)
4. [Health Information: ADHD and Hyperactivity](#)
5. [Health Information: ADHD Medicines: Suicide Warning for Strattera](#)
6. [Health Information: ADHD Myths and Facts](#)
7. [Health Information: Impulsivity and Inattention](#)
8. [Health Information: Other Conditions With Similar Symptoms](#)
9. [Health Information: Social Skills Training](#)
10. [Health Information: Tests for Other Disorders](#)
11. [Lexicomp: Attention Deficit Hyperactivity Disorder \(ADHD\)](#)
12. [Lexicomp: Attention Deficit Hyperactivity Disorder \(ADHD\) Discharge Instructions](#)
13. [Lexicomp: Medicines for Attention Deficit Hyperactivity Disorder \(ADHD\)](#)

## **Scope**

### **Disease/Condition(s):**

Attention deficit and hyperactivity disorder (ADHD)

### **Clinical Specialty:**

Family Medicine, Neurology, Pediatrics, Psychiatry, and Psychology

### **Intended Users:**

Primary Care Physicians, Advanced Practice Providers, Psychiatrists, Psychologists

### **CPG objective(s):**

To provide evidence-based recommendations for the effective diagnosis and treatment of adult patients with ADHD.

### **Target Population:**

Adult patients (age 18 years or older).

### **Major Outcomes Considered:**

1. Incidence of comorbid disorders
2. Effectiveness of treatment
3. Adverse effects of medication

## Methodology

**Methods Used to Collect/Select the Evidence:** Evidence was selected using hand searches of published literature and electronic databases.

**Methods Used to Assess the Quality and Strength of the Evidence and Recommendations:** Recommendations developed during the workgroup meetings used the modified Grading of Recommendations Assessment, Development and Evaluation (GRADE) developed by the American Heart Association and American College of Cardiology (**Figure 1**) to establish evidence grades for each piece of literature and/or recommendation.

**Rating Scheme for the Strength of the Evidence and Recommendations:** See [Appendix A](#).

**Methods Used to Formulate the Recommendations:** Recommendations developed by external organizations were adopted while others were developed via group consensus through discussion of the literature evidence and expert experiences.


## Introduction

Attention Deficit Hyperactivity Disorder, originally thought to occur just in childhood, is now widely understood as persisting into adulthood. Between 50 to 65 percent of adults diagnosed with childhood ADHD will continue to have symptoms of inattention, distractibility and impulsivity causing functional impairment as adults. In addition, adults who were never diagnosed as children may present with a complicated array of behavioral, legal and functional problems requesting diagnosis and treatment.

This guideline is designed to provide primary care clinicians with a structure, tools and referral criteria for diagnosis and treatment of adults 18 and over with symptoms typical of ADHD.

# Recommendations

## Adult ADHD Algorithm (ages 18 years or older)



**UWHealth**

Last reviewed/revised: 10/2014  
For revisions, contact CCKM.  
ADHD- Adult – Ambulatory Guideline

## 1. PRESENTATION AND SCREENING

Adults with potential ADHD may present with a self-diagnosis, at the suggestion of a family member, friend, employer or therapist or with other behavioral or psychological problems. (Class I, LOE B) There may or may not be a previous childhood or adult diagnosis of ADHD.

Adult ADHD is commonly characterized by poor executive functioning. Indicators of ADHD and screening symptoms include:

- Inattention
- Restlessness
- Forgetfulness
- Disorganization
- Impulsive behaviors/often impatient
- Poor planning
- Increased risk of driving and other accidents
- Family and relationship difficulties
- Difficulties with parenting

High risk behaviors, failed relationships, legal difficulties, substance abuse and recurrent job loss are common. Physical hyperactivity diminishes in severity with age, but inattentive symptoms become more prominent and may be perceived as incompetence. Some adults compensate by finding a spouse / partner who organizes them or a job which is very active, highly absorbing or stimulating.

## 2. CLINICAL ASSESSMENT

Evaluation of adults presenting with ADHD symptoms typically requires at least two visits. As well as allowing for a thorough evaluation, two visits allows the clinician to assess motivation for follow up, persistence of symptoms and dysfunction and likelihood for alternative diagnoses. The following components of a complete evaluation are considered during both visits (Class I, LOE C):

- review and corroboration of current symptoms and dysfunction
- determination of a childhood onset
- evaluation for comorbid and /or mimicking psychiatric problems, medical disorders or substance abuse.

### First Visit

#### A. Review Current Symptoms and Functional Impairment (Class I, LOE C)

- DSM-5 diagnostic criteria for ADHD should be used and followed. A validated adult ADHD assessment tool (such as the [Adult ADHD Self-Report Scale \(ASRS-v1.1\) Symptom Checklist](#)) may be used to adjunctively evaluate an adult patient.
- Adults may present with distractibility, impulsiveness and poor executive functioning. A variety of psychiatric or lifestyle conditions need to be considered when these symptoms are present.

**B. Establish Onset** *(Class I, LOE C)*

- ADHD is a neurodevelopmental disorder that may persist into adulthood.
- In order to meet diagnostic criteria, symptoms and functional impairment need to have been present in patients prior to age 12.

**C. Perform Medical Evaluation** *(Class I, LOE C)*

- Screen for medical, psychiatric or substance abuse issues which could explain or exacerbate symptoms of ADHD. *(Class I, LOE C)*
- Screen for medical and psychological conditions which would influence choice of medication. When considering a stimulant in an adult with risk factors for cardiac disease, the provider should consider a cardiovascular evaluation before initiating therapy. *(Class I, LOE C)*
- Establish baseline vital signs: weight, blood pressure, pulse. *(Class I, LOE C)*
- Laboratory testing should be limited to areas of concern. *(Class I, LOE C)*

**D. Evaluate for Psychiatric or Lifestyle Conditions**

- Adults may present with distractibility, impulsiveness and poor executive functioning. A variety of psychiatric or lifestyle conditions need to be considered when these symptoms are present. *(Class I, LOE C)*

**GATHER ADDITIONAL INFORMATION**

**A. Corroborate Childhood Onset and Impairment**

Childhood history can be gathered by review of medical records, review of report cards or other academic materials, and interview with parents or close family member either in person or via a phone call. High activity patterns, difficult temperament, and frequent accidents or risk taking behavior are common childhood characteristics. Review of academic background should reveal areas of impairment or concern. Look for drop outs, failures, learning disability, special evaluations or classes, suspensions / expulsions, and focused problems in areas such as reading, writing, penmanship or math. *(Class I, LOE C)*

Review of report cards often indicates behavior problems, lack of expected achievement, incomplete work, or inadequate effort. If there is no objective evidence of childhood symptoms and impairment, the diagnosis of adult ADHD should be reconsidered.

**B. Review Family Psychiatric History**

It is common to have a positive family psychiatric history. Inquire particularly about learning disabilities, behavior problems, legal difficulties, ADHD, and substance abuse. *(Class I, LOE B)*

**CONSIDER COMORBID OR ALTERNATIVE PSYCHIATRIC DIAGNOSIS**

*(Class I, LOE B)*

Psychiatric disorders can cause inattentive symptoms or can influence the course of treatment. Presence of another psychiatric diagnosis does not preclude a diagnosis of adult ADHD but it does make diagnosis and treatment more confusing. Significant

physical, verbal or emotional abuse / neglect can contribute to symptoms characteristic of ADHD. Depression, Post-Traumatic Stress Disorder (PTSD), bipolar disorder, anxiety disorder, personality disorders, substance abuse and other psychiatric disorders should be considered as a part of the evaluation. For recommendations related to assessment for depression, reference the [UW Health Depression – Adult – Guideline](#).

Patients whose psychiatric status is unclear should be referred to a mental health provider. Patients with active substance abuse should be referred to a substance use treatment program. For assessment of tobacco and alcohol use, reference the [UW Health Tobacco – Pediatric/Adult – Inpatient/Ambulatory Guideline](#) or [UW Health Alcohol – Pediatric/Adult – Ambulatory Guideline](#). Consider evaluation for drug-seeking behavior with multiple pharmacies or prescribing providers using the [Wisconsin Prescription Drug Monitoring Program](#).

It is important to identify comorbid disorders because they can mimic ADHD.

- a. Comorbid or alternative psychiatric conditions should be addressed prior to starting treatment for ADHD.
- b. Certain medical conditions (liver disease, seizures, hypertension, glaucoma) are relative contraindications to certain ADHD medications.

### **CONSIDER REFERRAL** *(Class I, LOE C)*

Referral to psychiatrists and additional providers is always at the discretion of the provider. There are several presentations and co-conditions for which referral is recommended:

1. Extreme dysfunction
2. Suicidality or homicidality
3. Substance abuse or dependence
4. Psychosis
5. Extreme psychosocial stressors
6. Previous treatment failures
7. Atypical presentation – if presentation as brand new symptoms this is not ADHD, even if not diagnosed as a child the symptoms must concur

### **3. ESTABLISH DIAGNOSIS**

To diagnose ADHD, the clinician should determine that DSM-5 criteria have been met. *(Class I, LOE B)*

#### **DSM-5 Diagnostic Criteria**

- A. A persistent pattern of inattention and/or hyperactivity-impulsivity that interferes with functioning or development, as characterized by (1) and/or (2):
  1. **Inattention:** Six (or more) of the following symptoms have persisted for at least 6 months to a degree that is inconsistent with developmental level and that negatively impacts directly on social and academic/occupational activities:

**Note:** The symptoms are not solely a manifestation of oppositional behavior, defiance, hostility, or failure to understand tasks or instructions. **For older adolescents and adults (age 17 and older), at least five symptoms are required.**

- a. Often fails to give close attention to details or makes careless mistakes in schoolwork, at work, or during other activities (e.g., overlooks or misses details, work is inaccurate).
- b. Often has difficulty sustaining attention in tasks or play activities (e.g., has difficulty remaining focused during lectures, conversations, or lengthy reading).
- c. Often does not seem to listen when spoken to directly (e.g., mind seems elsewhere, even in the absence of any obvious distraction).
- d. Often does not follow through on instructions and fails to finish schoolwork, chores, or duties in the workplace (e.g., starts tasks but quickly loses focus and is easily sidetracked).
- e. Often has difficulty organizing tasks and activities (e.g., difficulty managing sequential tasks; difficulty keeping materials and belongings in order; messy, disorganized work; has poor time management; fails to meet deadlines).
- f. Often avoids, dislikes, or is reluctant to engage in tasks that require sustained mental effort (e.g., schoolwork or homework; for older adolescents and adults, preparing reports, completing forms, reviewing lengthy papers).
- g. Often loses things necessary for tasks or activities (e.g., school materials, pencils, books, tools, wallets, keys, paperwork, eyeglasses, mobile telephones).
- h. Is often easily distracted by extraneous stimuli (for older adolescents and adults, may include unrelated thoughts).
- i. Is often forgetful in daily activities (e.g., doing chores, running errands; for older adolescents and adults, returning calls, paying bills, keeping appointments).

2. **Hyperactivity and impulsivity:** Six (or more) of the following symptoms have persisted for at least 6 months to a degree that is inconsistent with developmental level and that negatively impacts directly on social and academic/occupational activities:

**Note:** The symptoms are not solely a manifestation of oppositional behavior, defiance, hostility, or a failure to understand tasks or instructions. For older adolescents and adults (age 17 and older), at least five symptoms are required.

- a. Often fidgets with or taps hands or feet or squirms in seat.
- b. Often leaves seat in situations when remaining seated is expected (e.g., leaves his or her place in the classroom, in the office or other workplace, or in other situations that require remaining in place).
- c. Often runs about or climbs in situations where it is inappropriate. (**Note:** In adolescents or adults, may be limited to feeling restless.)
- d. Often unable to play or engage in leisure activities quietly.

- e. Is often “on the go,” acting as if “driven by a motor” (e.g., is unable to be or uncomfortable being still for extended time, as in restaurants, meetings; may be experienced by others as being restless or difficult to keep up with).
  - f. Often talks excessively.
  - g. Often blurts out an answer before a question has been completed (e.g., completes people’s sentences; cannot wait for turn in conversation).
  - h. Often has difficulty waiting his or her turn (e.g., while waiting in line).
  - i. Often interrupts or intrudes on others (e.g., butts into conversations, games, or activities; may start using other people’s things without asking or receiving permission; for adolescents and adults, may intrude into or take over what others are doing).
- B. Several inattentive or hyperactive-impulsive symptoms were present prior to age 12 years.
  - C. Several inattentive or hyperactive-impulsive symptoms are present in two or more settings (e.g., at home, school, or work; with friends or relatives; in other activities).
  - D. There is clear evidence that the symptoms interfere with, or reduce the quality of, social, academic, or occupational functioning.
  - E. The symptoms do not occur exclusively during the course of schizophrenia or another psychotic disorder and are not better explained by another mental disorder (e.g., mood disorder, anxiety disorder, dissociative disorder, personality disorder, substance intoxication or withdrawal).

### DSM-5 Diagnosis

*Specify whether:*

**Combined presentation:** If both Criterion A1 (inattention) and Criterion A2 (hyperactivity-impulsivity) are met for the past 6 months.

**Predominantly inattentive presentation:** If Criterion A1 (inattention) is met but Criterion A2 (hyperactivity-impulsivity) is not met for the past 6 months.

**Predominately hyperactive/impulsive presentation:** If Criterion A2 (hyperactivity-impulsivity) is met and Criterion A1 (inattention) is not met for the past 6 months.

*Specify if:*

**In partial remission:** When full criteria were previously met, fewer than the full criteria have been met for the past 6 months, and the symptoms still result in impairment in social, academic, or occupational functioning.

*Specify current severity:*

**Mild:** Few, if any, symptoms in excess of those required to make the diagnosis are present, and symptoms result in no more than minor impairments in social or occupational functioning.

**Moderate:** Symptoms or functional impairment between “mild” and “severe” are present.

**Severe:** Many symptoms in excess of those required to make the diagnosis, or several symptoms that are particularly severe, are present, or the symptoms result in marked impairment in social or occupational functioning.

## 4. PROVIDE TREATMENT

*(Class I, LOE C unless otherwise indicated)*

1. Provide or offer referral regarding ADHD symptom management, and psycho-education or effective coping strategies for both the patient and family.

2. Follow medication treatment protocol and medication chart ([Appendix B](#) and [Appendix C](#)). (Class I, LOE A) Specific patient needs or wishes should be considered and therapy should be individualized.
3. Little data is available on the use of therapeutic stimulants in pregnancy, but currently they are not associated with major congenital malformations. Risks of discontinuation of therapy should be considered (e.g., driving, vocational responsibilities) along with the benefits for each individual patient. (Class IIb, LOE C)
4. Long term benefit should be assessed for each patient, especially those who continue treatment from a childhood diagnosis. A trial discontinuation of therapy can be considered as children age into adulthood to assess ongoing benefit of therapy.
5. In situations where there is increased risk of substance abuse or diversion, non stimulant preparations or slow release stimulants are preferred and can be used to initiate treatment. When crushed, slow release stimulants resemble immediate release preparations in terms of onset and effect.
6. Adults with ADHD who are also parents may benefit from therapy to assist them with parenting skills.
7. Consider vocational and/or educational accommodation.
8. For patients at high risk of substance abuse, consider establishing a drug contract or conducting periodic drug screens.
9. Adjuvant psychotherapy.

## 5. COMPLETE FOLLOW-UP CARE

Adults with a new diagnosis, uncontrolled symptoms or change in medication should be seen within 30 days by a clinician who can assess for side effects and adjust medication if needed. Monthly contacts or visits should be routine until functionality is significantly improved. Once functionality is improved, follow-up appointments every 3 to 6 months are recommended. Informants should be included, as available, in follow-up sessions. (Class I, LOE C)

At each follow-up visit (Class I, LOE C):

1. Review should specifically include diurnal variation in symptoms, as this informs recommendations for change in timing/formulation of the medications prescribed.
2. Review target symptoms, job/school performance, relationship issues.
3. Monitor for adherence to therapy, drug side effects/toxicity or signs of abuse/diversion. Also monitor vital signs to assess for increases in blood pressure and pulse.
4. Review impressions of informants.
5. Adjust therapy as needed.

Medications must be prescribed in accordance with [Wisconsin Chapter 961](#) for controlled substances:

1. Prescription must be written for legitimate medical indication.
2. Sign/date prescription on date of issue with:
  - a. Patient full name/address.

- b. Drug name, strength, dosage form, quantity, directions for use.
3. Up to 3 monthly prescriptions may be given to patients.
  - a. The date of issue (date of prescription is written) must be on all three prescriptions.
  - b. The prescriber writes “fill on or after XX/XX/XXXX” for two prescriptions to be filled at a later date.
  - c. A prescription for a CII controlled substance cannot be dispensed more than 60 days after the date of issue on the prescription order.

## **UW Health Implementation**

**Potential Benefits:** Appropriate assessment and treatment of adults with attention-deficit/hyperactivity disorder.

**Potential Harms:** Drug toxicity.

### **Qualifying Statements**

The listed practice parameters are developed to assist clinicians in psychiatric decision making. These parameters are not intended to define the standard of care, nor should they be deemed inclusive of all proper methods of care or exclusive of other methods of care directed at obtaining the desired results. The ultimate judgment regarding the care of a particular patient must be made by the clinician in light of all of the circumstances presented by the patient and his or her family, the diagnostic and treatment options available, and available resources.

### **Implementation Plan/Tools**

1. Guideline will be housed on U-Connect in a dedicated folder for CPGs.
2. Release of the guideline will be advertised in the Clinical Knowledge Management Corner within the Best Practice newsletter.
3. Links to this guideline will be updated and/or added in appropriate Health Link or equivalent tools, including:
  - ADD/ADHD [73]

### **Disclaimer**

CPGs are described to assist clinicians by providing a framework for the evaluation and treatment of patients. This Clinical Practice Guideline outlines the preferred approach for most patients. It is not intended to replace a clinician’s judgment or to establish a protocol for all patients. It is understood that some patients will not fit the clinical condition contemplated by a guideline and that a guideline will rarely establish the only appropriate approach to a problem.

## References

1. Meszaros A, Czobor P, Balint S, Komlosi S, Simon V, Bitter I. Pharmacotherapy of adult attention deficit hyperactivity disorder (ADHD): a meta-analysis. *Int J Neuropsychopharmacol*. 2009;12(8): 1137-1147.
2. Faraone, SV, Glatt SJ. A comparison of the efficacy of medications for adult attention-deficit/hyperactivity disorder using meta-analysis of effect sizes. *J Clin Psychiatry* 2010; 71(6): 754-763.
3. Weiss N. Assessment and treatment of ADHD in adults. *Psychiatric Annals*. 2011; 41(1): 23-31. *Pediatrics* 2011; 128: 1007-1022.
4. Greenhill, Laurence, Pliszka, Steven, et al., Practice Parameter for the Use of Stimulant Medications in the Treatment of Children, Adolescents, and Adults; *J Am Acad of Child Adoles Psychiatry*. 2002; 41 (2 Supplement): 26S – 49S.
5. Kessler, Ronald; Adler, Lenard; et al, Patterns and Predictors of Attention-Deficit/Hyperactivity Disorder Persistence into Adulthood: Results from the National Comorbidity Survey Replication. *Biol Psychiatry* 2005; 57: 1442-1451.
6. Kessler, Ronald; Adler, Lenard; et al, The World Health Organization Adult ADHD Self-Report Scale (ASRS): a Short Screening Scale for Use in the General Population. *Psychological Medicine*, 2005; 35: 245-256.
7. McGough, James J and Russell Barkley, Diagnostic Controversies in Adult Attention Deficit Hyperactivity Disorder. *Am J Psychiatry* 2004; 161:1948-1956.
8. Montano, Brendan, Diagnosis and Treatment of ADHD in Adults in Primary Care. *J Clin Psychiatry* 2004; 65: (suppl 3): 18-21.
9. Searight, Russell, Burke, John and Fred Rottnek, Adult ADHD: Evaluation and Treatment in Family Medicine. *Am Fam Physician* 2000; 62: 2077-86.
10. Weiss, Margaret and Candice Murray, Assessment and Management of Attention-Deficit Hyperactivity Disorder in Adults. *SMAJ*; 168: 715-22.
11. Meijer WM, Faber A, van den Ban E, and Tobi H. Current Issues Around the Pharmacotherapy of ADHD in Children and Adults. *Pharm World Sci* 2009; 31:509-16.
12. Nutt DJ, Fone K, Asherson P, Bramble D, et al. Evidence-based guidelines for management of attention deficit/hyperactivity disorder in adolescents in transition to adult services and in adults: Recommendations from the British Association for Psychopharmacology. *J Psychopharmacol* 2007; 21: 10-41.
13. National Collaborating Center for Mental Health, Attention deficit hyperactivity disorder. Diagnosis and management of ADHD in children, young people and adults. London (UK): National Institute for Health and Clinical Excellence (NICE) 2008 Sep. 59 p (Clinical guideline; no. 72).
14. Surman Craig BH. ADHD in Adults. *Medscape* 2005 Weiss Nicholas. Assessment and Treatment of ADHD in Adults. (*Psychiatric Annals*) 2011 Jan: Vol 41:1.
15. Freeman, MP. ADHD and pregnancy. *Am J Psychiatry* 2014; 171:723-728.
16. Volkow, ND, Swanson, JM. Clinical practice: Adult attention deficit-hyperactivity disorder. *New England Journal of Medicine* 2013;369:1935-1944.


# Appendix A

Figure 1. AHA/ACC Modified GRADE Grading Scheme

		SIZE OF TREATMENT EFFECT												
		CLASS I	CLASS IIa	CLASS IIb	CLASS III <i>No Benefit</i> or CLASS III <i>Harm</i>									
		<b>Benefit &gt;&gt;&gt; Risk</b> Procedure/Treatment <b>SHOULD</b> be performed/administered	<b>Benefit &gt;&gt; Risk</b> Additional studies with <i>focused objectives needed</i> <b>IT IS REASONABLE</b> to perform procedure/administer treatment	<b>Benefit ≥ Risk</b> Additional studies with <i>broad objectives needed; additional registry data would be helpful</i> Procedure/Treatment <b>MAY BE CONSIDERED</b>	<table border="1"> <thead> <tr> <th></th> <th>Procedure/ Test</th> <th>Treatment</th> </tr> </thead> <tbody> <tr> <td>COR III: No benefit</td> <td>Not Helpful</td> <td>No Proven Benefit</td> </tr> <tr> <td>COR III: Harm</td> <td>Excess Cost w/o Benefit or Harmful</td> <td>Harmful to Patients</td> </tr> </tbody> </table>		Procedure/ Test	Treatment	COR III: No benefit	Not Helpful	No Proven Benefit	COR III: Harm	Excess Cost w/o Benefit or Harmful	Harmful to Patients
	Procedure/ Test	Treatment												
COR III: No benefit	Not Helpful	No Proven Benefit												
COR III: Harm	Excess Cost w/o Benefit or Harmful	Harmful to Patients												
ESTIMATE OF CERTAINTY (PRECISION) OF TREATMENT EFFECT	<b>LEVEL A</b> Multiple populations evaluated* Data derived from multiple randomized clinical trials or meta-analyses	<ul style="list-style-type: none"> <li>Recommendation that procedure or treatment is useful/effective</li> <li>Sufficient evidence from multiple randomized trials or meta-analyses</li> </ul>	<ul style="list-style-type: none"> <li>Recommendation in favor of treatment or procedure being useful/effective</li> <li>Some conflicting evidence from multiple randomized trials or meta-analyses</li> </ul>	<ul style="list-style-type: none"> <li>Recommendation's usefulness/efficacy less well established</li> <li>Greater conflicting evidence from multiple randomized trials or meta-analyses</li> </ul>	<ul style="list-style-type: none"> <li>Recommendation that procedure or treatment is not useful/effective and may be harmful</li> <li>Sufficient evidence from multiple randomized trials or meta-analyses</li> </ul>									
	<b>LEVEL B</b> Limited populations evaluated* Data derived from a single randomized trial or nonrandomized studies	<ul style="list-style-type: none"> <li>Recommendation that procedure or treatment is useful/effective</li> <li>Evidence from single randomized trial or nonrandomized studies</li> </ul>	<ul style="list-style-type: none"> <li>Recommendation in favor of treatment or procedure being useful/effective</li> <li>Some conflicting evidence from single randomized trial or nonrandomized studies</li> </ul>	<ul style="list-style-type: none"> <li>Recommendation's usefulness/efficacy less well established</li> <li>Greater conflicting evidence from single randomized trial or nonrandomized studies</li> </ul>	<ul style="list-style-type: none"> <li>Recommendation that procedure or treatment is not useful/effective and may be harmful</li> <li>Evidence from single randomized trial or nonrandomized studies</li> </ul>									
	<b>LEVEL C</b> Very limited populations evaluated* Only consensus opinion of experts, case studies, or standard of care	<ul style="list-style-type: none"> <li>Recommendation that procedure or treatment is useful/effective</li> <li>Only expert opinion, case studies, or standard of care</li> </ul>	<ul style="list-style-type: none"> <li>Recommendation in favor of treatment or procedure being useful/effective</li> <li>Only diverging expert opinion, case studies, or standard of care</li> </ul>	<ul style="list-style-type: none"> <li>Recommendation's usefulness/efficacy less well established</li> <li>Only diverging expert opinion, case studies, or standard of care</li> </ul>	<ul style="list-style-type: none"> <li>Recommendation that procedure or treatment is not useful/effective and may be harmful</li> <li>Only expert opinion, case studies, or standard of care</li> </ul>									

# Appendix B

## Adult ADHD Medication Algorithm



**UWHealth**  
Center for Clinical Knowledge Management

Last revised/reviewed: 10/2014  
ADHD- Adult – Ambulatory Guideline  
Contact CCKM for revisions.

## Appendix C

### Medications for Treatment of Attention-Deficit/Hyperactivity Disorder

#### GENERAL CONSIDERATIONS FOR STIMULANTS

<ul style="list-style-type: none"> <li>• Consider cardiac risk factors before initiating therapy</li> <li>• Use cautiously if history of tics</li> <li>• Give with/after food and swallow whole with liquids</li> <li>• Longer-acting stimulants may have greater problematic effects on evening appetite and sleep</li> <li>• Use cautiously if history of substance abuse or diversion concern</li> </ul>	<ul style="list-style-type: none"> <li>• Monitor patient weight and vital signs</li> <li>• Pellet/beaded capsule formulation may be opened and sprinkled on soft food</li> <li>• Nonabsorbable tablet shell may be seen in stool (Concerta)</li> <li>• Consider cardiovascular evaluation before initiating therapy</li> </ul>
---	--

#### Methylphenidate Products

	Product Names	Strengths Available	Duration of Action	Usual Dosing Adult Titration Dose (titrate every 7 days, unless otherwise indicated)	Maximum Daily Dose
Short acting	methylphenidate tab <sup>^*</sup> (Ritalin)	5,10, 20 mg tab	≤ 4 hours	5-20 mg given 2-3 times daily Titrate by 5-10 mg every 7-14 days	FDA: 60 mg Off label: 100 mg if over 50 kg
	methylphenidate <sup>^*</sup> (Methylin) (equivalent to Ritalin)	2.5, 5, 10mg chew tab 5 mg/5mL, 10mg/5mL solution	≤ 4 hours	5–20 mg given 2-3 times daily Titrate by 5-10 mg every 7-14 days	FDA: 60 mg Off label: 100 mg if over 50 kg
Intermediate acting  4-6 hours	methylphenidate SR tab <sup>^*</sup> (Ritalin SR) Medadate ER and generics rated AB equivalent	20 mg tab	4 – 6 hours	20–60 mg (divided in 1-2 doses/day) (20-40 mg in morning, 20 mg in early afternoon) Titrate by 20 mg/day	FDA: 60 mg Off label: 100 mg if over 50 kg
	methylphenidate <sup>^*</sup> (Methylin ER) (equivalent to Ritalin SR)	10,20 mg tablet	4 – 6 hours	10-60 mg daily	FDA: 60 mg Off label: 100 mg if over 50 kg
	Methylphenidate tab <sup>^*</sup> (Metadate ER)	20 mg tablet	4 – 6 hours	20-60 mg daily (divided in 1-2 doses/day)	FDA: 60 mg Off label: 100 mg if over 50 kg
	dexmethylphenidate <sup>^*</sup> (Focalin) cap	2.5, 5, 10 mg tab	4 – 6 hours	2.5–10 mg given twice daily at least 4 hours apart	FDA: 20 mg Off label: 50 mg

	Product Names	Strengths Available	Duration of Action	Usual Dosing Adult Titration Dose (titrate every 7 days, unless otherwise indicated)	Maximum Daily Dose
Intermediate acting 6-8 hours	methylphenidate*^ (Metadate CD) cap (bimodal release with 30% immediate release and 70% delayed release)	10, 20, 30, 40, 50, 60 mg capsule	6 – 8 hours	10-60mg daily Titration 10-20 mg	FDA: 60 mg Off label: 100 mg if over 50 kg
	methylphenidate ER*^§ (Ritalin LA) cap (bimodal release with 50% rapid onset and 50% delayed release)	10, 20, 30, 40 mg capsule	6 – 8 hours	20-60mg daily	FDA: 60 mg Off label: 100 mg if over 50 kg
Long acting	dexmethylphenidate*^§ (Focalin XR) (bimodal release with 50% immediate release and 50% delayed release)	5, 10, 15, 20, 25, 30, 35, 40 mg capsule	10 - 12 hours 5-20 mg once daily	5–40 mg daily	FDA: 40 mg Off label: 50 mg
	methylphenidate ^ (Daytrana) patch apply to hip for 9 hours	10, 15, 20, 30 mg patch	12 hours (with 2 -3 hour delay)	10-30mg patch daily Titrate by next highest strength patch	FDA: 30 mg
	Methylphenidate*^§ (Concerta) tab (bimodal release with immediate onset and delayed release)	18, 27, 36, 54 mg tab	10 hours	18-54mg once daily (titrate by 18 mg)	FDA: 54 mg for children, 72 mg for adolescents and adults Off label: 90 mg adolescents (>40 kg)
^ FDA approved for treatment of ADHD, * Generic product, §Oral long acting methylphenidate products have immediate release and extended release components.					



Center for Clinical Knowledge Management

Last revised/reviewed: 10/2014

ADHD – Adult – Ambulatory Guideline

Contact CCKM for revisions.

## Amphetamine Products

	Product Names	Strengths Available	Duration of action	Usual Dosing (titrate every 7 days, unless otherwise noted)	Maximum Dose
Short acting	Dextroamphetamine*	5, 10 mg tablet 1 mg/mL solution	4-6 hours	2.5 -15 mg two to three times Daily Titration 5 mg/week	FDA: 40 mg Off label: 60 mg (>50 kg)
Intermediate acting	dextroamphetamine capsule SR*§ (Dexedrine spansules) (bimodal release with 50% immediate release and 50% delayed release)	5, 10, 15 mg capsule	6-8 hours	5-15mg 2 timestwice daily Titration 5 mg	FDA: 40 mg Off label: 60 mg (>50 kg)
	amphetamine mixed salts tab ^combo* (Adderall) *	5, 7.5, 10, 12.5, 15, 20, 30 mg tab	5-8 hours	52.5-30mg 1-2 times once or twice daily Titration 2.5-5 mg once or twice daily	FDA: 40 mg Off label: 40 mg (≤ 50kg), 60 mg (>50 kg)
Long acting	amphetamine mixed salts capsule^* combo (Adderall XR)*§ (bimodal release with 50% immediate release and 50% delayed release)	5, 10, 15, 20, 25, 30 capsule	10 hours	10-30mg once daily Titration 5-10 mg	FDA: 30 mg Off-label: 30 mg (≤ 50kg), 60 mg (>50 kg)
	lisdexamfetamine (Vyvanse) capsule^	20, 30, 40, 50, 60, 70 mg capsule	10-12 hours	20-70mg once daily Titration 10-20 mg daily	FDA: 70 mg

^ FDA approved for treatment of ADHD, \* Generic product, §Oral long acting methylphenidate products have immediate release and extended release components.



Center for Clinical  
Knowledge Management

Last revised/reviewed: 10/2014  
ADHD – Adult – Ambulatory Guideline  
Contact CCKM for revisions.

## GENERAL CONSIDERATIONS FOR NON-STIMULANTS

<ul style="list-style-type: none"> <li>• May be used in cases of history of tics worsening from stimulants</li> <li>• Avoid bupropion if history of seizure or eating disorders</li> <li>• Monitor closely for behavioral side effects including suicidal ideation with atomoxetine, tricyclics, and bupropion as identified in FDA Black Box warning for anti-depressants</li> </ul>	<ul style="list-style-type: none"> <li>• Give with/after food and swallow whole with liquids</li> <li>• Medication of choice if concern about abuse or diversion</li> <li>• Consider cardiovascular risk factors before initiating tricyclic therapy and evaluate further if needed</li> <li>• Consider initiation with lower doses to improve tolerability</li> <li>• Guanfacine and clonidine may be used as adjunctive therapy with stimulants.</li> </ul>
---	---

### Non-Stimulant Products

	Product Names	Strengths Available	Duration of Action	Usual Dosing	Maximum Dosing
Anti-depressants	nortriptyline* (Pamelor, Aventyl)	10, 25, 50, 75 mg capsule 10 mg/5 mL solution	8-24 hours	0.5 mg/kg/day (May divide dose to 2-3 times daily)	2 mg/kg or 100 mg (whichever is lowest)
	bupropion* (Wellbutrin)	75, 100 mg tab	4-5 hours	3 -6 mg/kg/day (or 150 mg – 300 mg, whichever is lowest) Divide into 2 or 3 daily doses	6 mg/kg/day (or 300 mg) Whichever is lowest Divide into 2 or 3 daily dose
	bupropion SR* (Wellbutrin SR)	100, 150, 200 mg tab	12 hours	3 -6 mg/kg/day (or 150 mg – 300 mg, whichever is lowest) Divide into 2 daily doses.	6 mg/kg/day (or 300 mg) whichever is lowest Divide into 2 daily doses.
	bupropion XL* (Wellbutrin XL)	150, 300 mg tab	24 hours	3 -6 mg/kg/day (or 150 mg – 300 mg, whichever is lowest)	6 mg/kg/day (or 300 mg) whichever is lowest
Alpha-agonists	clonidine tab ER <sup>^</sup> (Kapvay)	0.1, 0.2 mg tab	At least 10-12 hours	0.1-0.4 mg/day Titration: 0.1 mg every 7 days	0.4 mg/day
	clonidine* (Catapres)	0.1, 0.2, 0.3 mg tab	At least 4-6 hours	0.05 mg at bedtime; 0.1 mg (≥ 45 kg) Titrate by 0.05 mg (<45 kg) or 0.1 mg (≥ 45 kg) increments to twice daily, three times daily, four times daily	0.4 mg (>45 kg)

	Product Names	Strengths Available	Duration of Action	Usual Dosing	Maximum Dosing
Alpha-agonists	guanfacine* (Tenex)	1, 2 mg tab	6-8 hours	0.5 mg at bedtime (<45 kg), 1 mg at bedtime (≥ 45 kg) Titrate by 0.5 mg (<45 kg) or 1 mg (≥ 45 kg) increments to twice daily, three times daily, four times daily	0.4 mg (>45 kg)
	guanfacine tab ER^* (Intuniv)	1, 2, 3, 4 mg tabs	At least 10-12 hours	0.05-0.12 mg/kg daily (or 1-4 mg once daily) Titration: 1 mg every 7 days	4 mg/day
Norepinephrine reuptake inhibitor	atomoxetine^ (Strattera) capsule	10, 18, 25, 40, 60, 80, 100 mg capsule	At least 10-12 hours	Initial dose )40 mg/day) After ≥ 3 days (increase to 80 mg daily)	FDA: 100 mg/day

\*Generic product

^ FDA Approved



Last revised/reviewed: 10/2014

ADHD – Adult – Ambulatory Guideline  
Contact CCKM for revisions.

### Potential Harms: Side Effects of Pharmacotherapy

- *Stimulants*: The most common side effects include appetite decrease, weight loss, insomnia, or headache. Less common side effects include tics and emotional lability/irritability, liver toxicity, hypertension, cardiac arrhythmia and psychosis.
- *Atomoxetine*: Side effects of atomoxetine that occurred more often than those with placebo include gastrointestinal distress, sedation, and decreased appetite.
- The U.S. Food and Drug Administration (FDA) and its Pediatric Advisory Committee have reviewed data regarding psychiatric adverse events to medications for the treatment of attention deficit/hyperactivity disorder (ADHD). For each agent examined (all stimulants, atomoxetine, and modafinil), there were reports of rare events of psychotic symptoms, specifically involving visual and tactile hallucinations of insects. Symptoms of aggression, suicidality (but no completed suicides), and cardiovascular issues were also reported.
- *Bupropion* may cause mild insomnia or loss of appetite. The highest recommended dose of bupropion is 450 mg. Higher doses may increase the risk of seizure.
- *Tricyclic Antidepressants (TCAs)* such as nortriptyline - frequently cause anticholinergic side effects such as dry mouth, sedation, constipation, changes in vision, or tachycardia. Among the TCAs, desipramine should be used with extreme caution in children and adolescents because there have been reports of sudden death. For TCAs electrocardiography should be considered for patients at risk and be performed at baseline and after each dose increase. Once the patient is on a stable dose of the TCA, a plasma level should be obtained to ensure the level is not in the toxic range.
- *Alpha-agonists*: Side effects of alpha-agonists include sedation, dizziness, and possible hypotension. Abrupt discontinuations of alpha-agonist are to be avoided.
- *Combinations of Medications*: There have been four deaths reported to the FDA of children taking a combination of methylphenidate and clonidine, but there were many atypical aspects of these cases.

# Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist

## Instructions

*The questions on the back page are designed to stimulate dialogue between you and your patients and to help confirm if they may be suffering from the symptoms of attention-deficit/hyperactivity disorder (ADHD).*

Description: The Symptom Checklist is an instrument consisting of the eighteen DSM-IV-TR criteria. Six of the eighteen questions were found to be the most predictive of symptoms consistent with ADHD. These six questions are the basis for the ASRS v1.1 Screener and are also Part A of the Symptom Checklist. Part B of the Symptom Checklist contains the remaining twelve questions.

### Instructions:

#### Symptoms

1. Ask the patient to complete both Part A and Part B of the Symptom Checklist by marking an X in the box that most closely represents the frequency of occurrence of each of the symptoms.
2. Score Part A. If four or more marks appear in the darkly shaded boxes within Part A then the patient has symptoms highly consistent with ADHD in adults and further investigation is warranted.
3. The frequency scores on Part B provide additional cues and can serve as further probes into the patient's symptoms. Pay particular attention to marks appearing in the dark shaded boxes. The frequency-based response is more sensitive with certain questions. No total score or diagnostic likelihood is utilized for the twelve questions. It has been found that the six questions in Part A are the most predictive of the disorder and are best for use as a screening instrument.

#### Impairments

1. Review the entire Symptom Checklist with your patients and evaluate the level of impairment associated with the symptom.
2. Consider work/school, social and family settings.
3. Symptom frequency is often associated with symptom severity, therefore the Symptom Checklist may also aid in the assessment of impairments. If your patients have frequent symptoms, you may want to ask them to describe how these problems have affected the ability to work, take care of things at home, or get along with other people such as their spouse/significant other.

#### History

1. Assess the presence of these symptoms or similar symptoms in childhood. Adults who have ADHD need not have been formally diagnosed in childhood. In evaluating a patient's history, look for evidence of early-appearing and long-standing problems with attention or self-control. Some significant symptoms should have been present in childhood, but full symptomology is not necessary.

# Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist

Patient Name		Today's Date					
Please answer the questions below, rating yourself on each of the criteria shown using the scale on the right side of the page. As you answer each question, place an X in the box that best describes how you have felt and conducted yourself over the past 6 months. Please give this completed checklist to your healthcare professional to discuss during today's appointment.			Never	Rarely	Sometimes	Often	Very Often
1. How often do you have trouble wrapping up the final details of a project, once the challenging parts have been done?							
2. How often do you have difficulty getting things in order when you have to do a task that requires organization?							
3. How often do you have problems remembering appointments or obligations?							
4. When you have a task that requires a lot of thought, how often do you avoid or delay getting started?							
5. How often do you fidget or squirm with your hands or feet when you have to sit down for a long time?							
6. How often do you feel overly active and compelled to do things, like you were driven by a motor?							
<b>Part A</b>							
7. How often do you make careless mistakes when you have to work on a boring or difficult project?							
8. How often do you have difficulty keeping your attention when you are doing boring or repetitive work?							
9. How often do you have difficulty concentrating on what people say to you, even when they are speaking to you directly?							
10. How often do you misplace or have difficulty finding things at home or at work?							
11. How often are you distracted by activity or noise around you?							
12. How often do you leave your seat in meetings or other situations in which you are expected to remain seated?							
13. How often do you feel restless or fidgety?							
14. How often do you have difficulty unwinding and relaxing when you have time to yourself?							
15. How often do you find yourself talking too much when you are in social situations?							
16. When you're in a conversation, how often do you find yourself finishing the sentences of the people you are talking to, before they can finish them themselves?							
17. How often do you have difficulty waiting your turn in situations when turn taking is required?							
18. How often do you interrupt others when they are busy?							
<b>Part B</b>							

## The Value of Screening for Adults With ADHD

Research suggests that the symptoms of ADHD can persist into adulthood, having a significant impact on the relationships, careers, and even the personal safety of your patients who may suffer from it.<sup>1-4</sup> Because this disorder is often misunderstood, many people who have it do not receive appropriate treatment and, as a result, may never reach their full potential. Part of the problem is that it can be difficult to diagnose, particularly in adults.

The Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist was developed in conjunction with the World Health Organization (WHO), and the Workgroup on Adult ADHD that included the following team of psychiatrists and researchers:

- **Lenard Adler, MD**  
Associate Professor of Psychiatry and Neurology  
New York University Medical School
- **Ronald C. Kessler, PhD**  
Professor, Department of Health Care Policy  
Harvard Medical School
- **Thomas Spencer, MD**  
Associate Professor of Psychiatry  
Harvard Medical School

As a healthcare professional, you can use the ASRS v1.1 as a tool to help screen for ADHD in adult patients. Insights gained through this screening may suggest the need for a more in-depth clinician interview. The questions in the ASRS v1.1 are consistent with DSM-IV criteria and address the manifestations of ADHD symptoms in adults. Content of the questionnaire also reflects the importance that DSM-IV places on symptoms, impairments, and history for a correct diagnosis.<sup>4</sup>

The checklist takes about 5 minutes to complete and can provide information that is critical to supplement the diagnostic process.

### References:

1. Schweitzer JB, et al. *Med Clin North Am.* 2001;85(3):10-11, 757-777.
2. Barkley RA. *Attention Deficit Hyperactivity Disorder: A Handbook for Diagnosis and Treatment.* 2nd ed. 1998.
3. Biederman J, et al. *Am J Psychiatry.* 1993;150:1792-1798.
4. American Psychiatric Association: *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision.* Washington, DC, American Psychiatric Association. 2000: 85-93.